

## Problem Set 2

Due Friday, April 18.

1. (Warm-up, don't hand in.) Exercise 27.4-2 (Generic Preflow-Push Max. Flow Algorithm, p. 614).
2. Potential function analysis of the Edmonds-Karp algorithm.

At any point during the execution of the Edmonds-Karp algorithm:

- Define the *shortest path in-degree* of a vertex  $v$  to be the number of edges coming in to  $v$  that are on shortest paths from the source to  $v$  in the residual graph.  
Let  $deg_{sp}(v)$  denote  $v$ 's shortest-path in-degree.
- Let  $level(v)$  denote the distance of  $v$  from the source in the residual graph.
- Define potential function  $\Phi$  to be  $\sum_v level(v)|E| - deg_{sp}(v)$ .

(a) Show that this potential function increases with every round of the Edmonds-Karp algorithm. Conclude that the number of rounds is  $O(V^2E)$ .

(b) Modify the potential function to improve the bound to  $O(VE)$ .

3. Show that the generic Preflow-Push Algorithm can in fact perform:

- (a)  $\Omega(V^2)$  lift operations and  $\Omega(V^2)$  saturating pushes
- (b) (Extra Credit)  $\Omega(V^3)$  nonsaturating pushes.

4. For the maximum-flow problem, the input is a directed graph  $G = (V, E)$ , a source vertex  $s$  and a sink vertex  $t$ , and, for each edge  $e \in E$ , a capacity  $c(e) > 0$ .

For the *minimum-cost flow problem*, one is also given a *cost*  $w(e) \geq 0$  for each edge  $e \in E$ . The *cost* of a flow  $f$  on  $G$  is defined to be

$$\sum_{e:f(e)>0} f(e)w(e).$$

The goal is to find a maximum flow that is of minimum cost among all maximum flows.

In the residual graph  $G_f = (V, E_f)$  (as defined in the book for the maximum-flow problem), define the *cost*  $w_f(u, v)$  of each edge  $(u, v) \in E_f$  to be  $-w(v, u)$  if  $f(v, u) > 0$ , or  $w(u, v)$  otherwise.

(a) Prove that if the residual graph  $G_f$  for a flow  $f$  contains no negative cost cycle (with respect to the cost function  $w_f$ ) then  $f$  is a minimum-cost flow among flows of value  $|f|$ .

(b) Among all  $s \rightsquigarrow t$  paths in the residual graph, let  $p$  be a shortest path, with respect to the cost function  $w_f$ . Prove that augmenting the flow  $f$  by sending flow along  $p$  yields a flow  $f'$  that is of minimum-cost among all flows of value  $|f'|$ .

(c) Why does this imply that in a network with integer capacities, there is always an integer minimum-cost maximum flow?