

CS 130 : Computer Graphics

Lecture 17: Physics-Based Simulation

Tamar Shinar

Computer Science & Engineering

UC Riverside

Physically-based simulation is widely used for movies, games,...

solid simulation in Pixar's *Ratatouille*

Physically-based simulation is widely used for movies, games,...



solid simulation in Pixar's *Ratatouille*

Physically-based simulation is widely used for movies, games,...

fluid simulation in Pixar's *Ratatouille*

Physically-based simulation is widely used for movies, games,...



fluid simulation in Pixar's *Ratatouille*

Firestorm

Harry Potter and the Half Blood Prince

Industrial Light + Magic



Firestorm

Harry Potter and the Half Blood Prince

Industrial Light + Magic

Similar techniques are used in 3D interactive games

- but here the constraint is real-time simulation
- detailed simulation for movies happens 'off-line'

Some of the basic types of objects we're interested in simulating

- rigid bodies
- soft bodies
 - hair, cloth, deformable volumetric solids
- fluids
 - smoke, liquids, fire

Particles

Particle: basic dynamic object



Particle: basic dynamic object



mass

m

Particle: basic dynamic object



mass

m

3 dof

$$\vec{X} = (x, y, z)$$

Particle: basic dynamic object



mass

$$m$$

3 dof

$$\vec{X} = (x, y, z)$$

forces: e.g., gravity

$$\vec{F} = -m\vec{g}$$

Particle: basic dynamic object



Equations of motion:
Newton's 2nd Law

$$\vec{F} = m\vec{a}$$

Particle: basic dynamic object



Equations of motion:
Newton's 2nd Law


$$\vec{F} = m\vec{a}$$



$$\frac{d\vec{x}}{dt} = \vec{v}$$
$$m \frac{d\vec{v}}{dt} = \vec{F}$$

System of
ODEs

Deformable bodies

Connect a bunch of particles into a 1D line segment with springs






A Mass Spring Model for Hair Simulation

Selle, A., Lentine, M., G., and Fedkiw, R. ACM Transactions on Graphics SIGGRAPH 2008, ACM TOG 27, 64.1-64.11 (2008)


Connect a bunch of particles into a 2D mesh








Selle, A., Su, J., Irving, G., and Fedkiw, R. IEEE Transactions on Visualization and Graphics (TVCG) 15(2) 339-350



Connect a bunch of particles into a 3D mesh



tetrahedron








Deformable bodies: equations of motion

Equations of motion:
Newton's 2nd Law



$$\vec{F} = m\vec{a}$$

$$\frac{d\vec{x}}{dt} = \vec{v}$$

$$m\frac{d\vec{v}}{dt} = \vec{F}$$

System of
PDEs

contains spatial derivatives