

# physics-based animation

## **Physics-Based Animation**

**Euler's Method**  $A = \frac{m}{m} \begin{pmatrix} M & B \\ B^{T} & 0 \end{pmatrix} \begin{pmatrix} M & B \\ B^{T} & 0 \end{pmatrix} \begin{pmatrix} M & B \\ B^{T} & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} b \\ c \end{pmatrix}$ VITKIN SIGGRAPH 2001 COURSE NOTES L E Rana, R E IR MARA ; L, R nonsingular S = diag(L, RT)  $S^{-1}AS^{-\tau} = \begin{pmatrix} L^{-1} \\ R^{-\tau} \end{pmatrix} \begin{pmatrix} M & B \\ B^{\tau} & o \end{pmatrix} \begin{pmatrix} L^{-\tau} \\ R^{-\tau} \end{pmatrix}$  $= \begin{pmatrix} L^{-1}ML^{-T} & L^{-1}BR^{-1} \\ R^{-T}B^{T}L^{-T} & 0 \end{pmatrix}$  $\mathbf{x}(t + \Delta t) = \mathbf{x}(t) + \Delta t \, \mathbf{f}(\mathbf{x}, t)$  $= \begin{pmatrix} L^{-1}ML^{-1} & L^{-1}BR^{-1} \\ (L^{-1}BR^{-1})^{T} & 0 \end{pmatrix}$ WITKIN SIGGRAPH 2001 COURS Particle system  $S^{-1}AS^{-T}S^{T}(\frac{x}{2}) = S^{-1}(\frac{b}{2})$  $\begin{pmatrix} L^{-1} M L^{-T} & L^{-1} \partial R^{-1} \\ (L^{-1} \partial R^{-1})^T & O \end{pmatrix} \begin{pmatrix} U^{-1} \\ W \end{pmatrix} = \begin{pmatrix} L^{-1} & B \\ R^{-T} & C \end{pmatrix}$ G sys where  $\begin{pmatrix} U^{T} \\ W \end{pmatrix} = S^{T} \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} L^{T} \\ R \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} L^{T} \\ R y \end{pmatrix}$ apply\_fun Choose L, F R. st K(L'ML) and K(L'BR') < K(D) p->f += p->m \* F->G P(L-ML-T, L-188-1) 2 P(M,B) Numerical Methods **Mathematics Physics** and Algorithms




#### Guendelman et al. 2003



# Shinar et al. 2008


Hong et al. 2007




#### Clausen et al. 2013


#### Zhu et al. 2015

Patkar et al. 2013





World Space



Material Space

Chentanez et al., 2009


# Physics of Natural Phenomena

#### Newton's Second Law (F = ma)

The acceleration **a** of a body is parallel and directly proportional to the net force **F** acting on the body, is in the direction of the net force, and is inversely proportional to the mass **m** of the body.

#### Newton's Third Law (Action/ Reaction)


When a body exerts a force  $F_1$  on a second body, the second body simultaneously exerts a force  $F_2=-F_1$  on the first body. This means that  $F_1$  and  $F_2$  are equal in magnitude and opposite in direction.



[Wikipedia]

# Math of Natural Phenomena


#### **Ordinary Differential Equations**




$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, t)$$

x(t): a moving point.
f(x,t): x's velocity.


## Math of Natural Phenomena



# Numerical Solution of Diff. Eq.



## Data Structures and Algorithms



- I. Advance velocity  $\mathbf{v}^n \to \tilde{\mathbf{v}}^{n+\frac{1}{2}}$
- II. Apply collisions  $\mathbf{v}^n \to \mathbf{\hat{v}}^n$ ,  $\mathbf{\tilde{v}}^{n+\frac{1}{2}} \to \mathbf{\hat{v}}^{n+\frac{1}{2}}$
- III. Apply contact and constraint forces  $\hat{\mathbf{v}}^{n+\frac{1}{2}} \rightarrow \mathbf{v}^{n+\frac{1}{2}}$
- IV. Advance positions  $\mathbf{x}^n \to \mathbf{x}^{n+1}$  using  $\mathbf{v}^{n+\frac{1}{2}}$ ,  $\mathbf{\hat{v}}^n \to \overline{\mathbf{v}}^n$
- V. Advance velocity  $\overline{\mathbf{v}}^n \to \mathbf{v}^{n+1}$

# Particles




mass m $\mathbf{3} \operatorname{dof}$  $\vec{X} = (x, y, z)$ 



mass m**3 dof**  $\vec{X} = (x, y, z)$ 


forces: e.g., gravity

$$\vec{F} = -m\vec{g}$$



#### Equations of motion: Newton's 2nd Law

$$\vec{F} = m\vec{a}$$



#### Equations of motion: Newton's 2nd Law


$$\vec{F} = m\vec{a}$$

$$\frac{d\vec{x}}{dt} = \vec{v}$$
$$m\frac{d\vec{v}}{dt} = \vec{F}$$

System of ODEs

# Deformable bodies


# Connect a bunch of particles into a <u>ID line</u> segment with springs



#### **A Mass Spring Model for Hair Simulation**


Selle, A., Lentine, M., G., and Fedkiw, R. ACM Transactions on Graphics SIGGRAPH 2008, ACM TOG 27, 64.1-64.11 (2008)


#### Connect a bunch of particles into a <u>2D mesh</u>




Selle et al. 2007


#### Connect a bunch of particles into a <u>3D mesh</u>







#### Deformable bodies: equations of motion



#### Equations of motion: Newton's 2nd Law

$$\vec{F} = m\vec{a}$$

$$\frac{d\vec{x}}{dt} = \vec{v}$$

$$m\frac{d\vec{v}}{dt} = \vec{F}$$

$$\mathbf{PDEs}$$


$$\mathbf{V}$$


contains spatial derivatives

# Rigid bodies

**Rigid bodies** 

6 dofs forces and torques elastic collisions ODEs







 $(\vec{F},\vec{\tau})$ 

#### Rigid body phenomena

# stacking collisions, conta friction articulation, control

#### Articulated rigid bodies





Rachel Weinstein, Joey Teran and Ron Fedkiw


# Rigid body simulation

#### [Weinstein et al 2006]



#### Rigid and deformable solids coupled together...


### Fracture





[Molino et al. 2004]

## Contact and collision


frame 25: (1.04167)









Simultaneous resolution of contact, elastic deformation, articulation constraints










our rigid/deformable simulator in Pixar's WALL-E

Selle et al. 2007


## Fluid simulation

# In fluid simulation, we often use a grid-based representation



An implicit representation has certain advantages over an explicit representation

- naturally handles topological changes
- very easy to extend from 2D to 3D



#### Fluid equations of motion: Navier-Stokes equations

$$\vec{F} = m\vec{a}$$

$$\rho(\mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u}) = \mu \triangle \mathbf{u} - \nabla p + \mathbf{f}$$

Hong et al. 2007


**Two-way Coupled SPH and Particle Level Set Fluid Simulation** 

Losasso, F., Talton, J., Kwatra, N. and Fedkiw, R. IEEE TVCG 14, No. 4 (2008)

### Control of virtual character

#### [Shinar et al. 2008]





rigid/deformable simulator in Pixar's WALL-E