

# Homework 5

## CS 210

Question	Points	Score
1	16	
2	16	
3	20	
4	18	
5	20	
Total	90	

1. (Heath 6.3) For each of the following functions, what do the first- and second- order optimality conditions say about whether 0 is a minimizer on  $\mathbb{R}$ ?

- (a)  $f(x) = x^2$
- (b)  $f(x) = x^3$
- (c)  $f(x) = x^4$
- (d)  $f(x) = -x^4$

2. (Heath 6.4) Determine the critical points of each of the following functions and characterize each as a minimum, maximum, or inflection point. Also determine whether each function has a global minimum or maximum on  $\mathbb{R}$ .

- (a)  $f(x) = x^3 + 6x^2 - 15x + 2$
- (b)  $f(x) = 2x^3 - 25x^2 - 12x + 15$
- (c)  $f(x) = 3x^3 + 7x^2 - 15x - 3$
- (d)  $f(x) = x^2e^x$

3. (Heath 6.5) Determine the critical points of each of the following functions and characterize each as a minimum, maximum, or saddle point. Also determine whether each function has a global minimum or maximum on  $\mathbb{R}^2$ .

- (a)  $f(x, y) = x^2 - 4xy + y^2$
- (b)  $f(x, y) = x^4 - 4xy + y^4$
- (c)  $f(x, y) = 2x^3 - 3x^2 - 6xy(x - y - 1)$
- (d)  $f(x, y) = (x - y)^4 + x^2 - y^2 - 2x + 2y + 1$

4. (Heath 6.8) Consider the function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  defined by

$$f(\mathbf{x}) = \frac{1}{2}(x_1^2 - x_2)^2 + \frac{1}{2}(1 - x_1)^2.$$

- (a) At what point does  $f$  attain a minimum?
- (b) Perform one iteration of Newton's method for minimizing  $f$  using as starting point  $\mathbf{x}_0 = (2, 2)^T$ .
- (c) Explain whether this is a good or bad step and in what sense.

5. (Heath 6.9) Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be given by

$$f(\mathbf{x}) = \frac{1}{2} \mathbf{x}^T A \mathbf{x} - \mathbf{x}^T \mathbf{b} + c$$

where  $A$  is an  $n \times n$  symmetric positive definite matrix,  $\mathbf{b}$  is an  $n$ -vector, and  $c$  is a scalar.

- (a) Show that Newton's method for minimizing this function converges in one iteration from any starting point  $\mathbf{x}_0$ .
- (b) If the steepest descent method is used on this problem, what happens if the starting value  $\mathbf{x}_0$  is such that  $\mathbf{x}_0 - \mathbf{x}^*$  is an eigenvector of  $A$ , where  $\mathbf{x}^*$  is the solution?